



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification ⁵ : C12P 17/14, C07D 277/34 C07D 263/44, 417/12, 413/12 A61K 31/425, 31/42</p>	A1	<p>(11) International Publication Number: WO 93/10254</p> <p>(43) International Publication Date: 27 May 1993 (27.05.93)</p>
<p>(21) International Application Number: PCT/GB92/02140</p> <p>(22) International Filing Date: 19 November 1992 (19.11.92)</p> <p>(30) Priority data: 9124513.4 19 November 1991 (19.11.91) GB</p> <p>(71) Applicant (for all designated States except US): SMITH-KLINE BEECHAM PLC. [GB/GB]; New Horizons Court, Brentford, Middlesex TW8 9EP (GB).</p> <p>(72) Inventors; and (75) Inventors/Applicants (for US only) : HINDLEY, Richard, Mark [GB/GB]; SmithKline Beecham Pharmaceuticals, Great Burgh, Yew Tree Bottom Road, Epsom, Surrey KT18 5XQ (GB). WORONIECKI, Stefan, Roland [GB/GB]; SmithKline Beecham Pharmaceuticals, Brockham Park, Betchworth, Surrey RH3 7AJ (GB).</p>		<p>(74) Agent: RUTTER, Keith; SmithKline Beecham, Corporate Patents, Great Burgh, Yew Tree Bottom Road, Epsom, Surrey KT18 5XQ (GB).</p> <p>(81) Designated States: AU, CA, JP, KR, US, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, SE).</p> <p>Published <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i></p>

(54) Title: PROCESS FOR THE PREPARATION OF PHARMACEUTICALLY ACTIVE THIAZOLIDINE OR OXAZOLIDINE COMPOUNDS BY A YEAST REDUCTASE



(I)



(II)

(57) Abstract

A process for preparing a compound of formula (I), or a tautomeric form thereof and/or a pharmaceutically acceptable salt thereof and/or a pharmaceutically acceptable solvate thereof, wherein: T represents a substituted or unsubstituted aryl group and T' is O or S; which process comprises, treating a compound of formula (II), or a tautomeric form thereof and/or a solvate thereof, wherein T and T' are as defined in relation to formula (I), with a microbial reductase obtained from an appropriate red yeast; and thereafter, as required, preparing a pharmaceutically acceptable salt and/or a pharmaceutically acceptable solvate of the compound of formula (I) or a tautomeric form thereof: compounds prepared by such a process and the use of such compounds in medicine.

FOR THE PURPOSES OF INFORMATION ONLY


Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	FR	France	MR	Mauritania
AU	Australia	GA	Gabon	MW	Malawi
BB	Barbados	GB	United Kingdom	NL	Netherlands
BE	Belgium	GN	Guinea	NO	Norway
BF	Burkina Faso	GR	Greece	NZ	New Zealand
BG	Bulgaria	HU	Hungary	PL	Poland
BJ	Benin	IE	Ireland	PT	Portugal
BR	Brazil	IT	Italy	RO	Romania
CA	Canada	JP	Japan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CC	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SK	Slovak Republic
CI	Côte d'Ivoire	LJ	Liechtenstein	SN	Senegal
CM	Cameroon	LK	Sri Lanka	SU	Soviet Union
CS	Czechoslovakia	LJ	Luxembourg	TD	Chad
CZ	Czech Republic	MC	Monaco	TC	Togo
DE	Germany	MG	Madagascar	UA	Ukraine
DK	Denmark	ML	Mali	US	United States of America
ES	Spain	MN	Mongolia	VN	Viet Nam
FI	Finland				

PROCESS FOR THE PREPARATION OF PHARMACEUTICALLY ACTIVE THIAZOLIDINE OR OXAZOLIDINE COMPOUNDS BY A YEAST REDUCTASE

This invention relates to a novel process for preparing certain substituted thiazolidinedione derivatives, to certain novel compounds, to pharmaceutical compositions comprising such compounds and to the use of such compounds in medicine.

European Patent Application, Publication Number 0 306228 discloses inter alia certain thiazolidinedione derivatives of formula (A):



(A)

or a tautomeric form thereof and/or a pharmaceutically acceptable salt thereof, and/or a pharmaceutically acceptable solvate thereof, wherein: A^a represents a substituted or unsubstituted aromatic heterocyclyl group;

R^a represents a hydrogen atom, an alkyl group, an acyl group, an aralkyl group, wherein the aryl moiety may be substituted or unsubstituted, or a substituted or unsubstituted aryl group;

R^b and R^c each represent hydrogen or R^b and R^c together represent a bond;

A^b represents a benzene ring having in total up to five substituents; and

n' represents an integer in the range of from 2 to 6.

EP 0306228 also discloses a process for converting compounds of formula (A), in particular a process for reducing the benzylidene double bond of compounds of formula (A) wherein R^b and R^c together represent a bond to provide compounds of formula (A) wherein R^b and R^c each represent hydrogen.


Particular reduction methods disclosed in EP 0306228 are catalytic reduction methods and metal/solvent reducing methods.

It has now been discovered that the reduction of the benzylidene double bond of the compound of formula (A) may be effected by a reductase enzyme from a microorganism, in particular from a yeast microorganism, and surprisingly that the said reduction proceeds in a structure specific manner. The reduction is also capable of proceeding in an enantioselective manner.

Certain of the compounds obtained in the process are novel and are indicated to show good blood-glucose lowering activity and are therefore of potential use in the treatment and/or prophylaxis of hyperglycaemia and are of particular use in the treatment of Type II diabetes.


These compounds are also indicated to be of potential use for the treatment and/or prophylaxis of other diseases including hyperlipidaemia, hypertension, cardiovascular disease and certain eating disorders.

Accordingly, the present invention provides a process for preparing a compound of formula (I):



(I)


or a tautomeric form thereof and/or a pharmaceutically acceptable salt thereof and/or a pharmaceutically acceptable solvate thereof, wherein: T represents a substituted or unsubstituted aryl group and T¹ is O or S; which process comprises, treating a compound of formula (II):



(II)

- or a tautomeric form thereof and/or a salt thereof and/or a solvate thereof, wherein T and T¹ are as defined in relation to formula (I), with a microbial reductase obtained from an appropriate red yeast; and
- 5 thereafter, as required, preparing a pharmaceutically acceptable salt and/or a pharmaceutically acceptable solvate of the compound of formula (I) or a tautomeric form thereof.


- Suitably T represents a moiety selected from the list consisting of (Ia),
- 10 (Ib), (Ic), (Id), (Ie), (If), (Ig), (Ih), (Ii), (Ij), (Ik), (Il), (Im), (In), (Io) and (Ip):



(Ia)

15


wherein A¹, A², R¹ and n are as defined in relation to formula (I) of EP 0306228;



(Ib)

20


wherein R², L¹, L² and L³ are as defined in relation to formula (I) of EP 0008203;



(Ic)


25

wherein R¹, R², R³, R⁴, R⁵, W and n are as defined in relation to formula (I) of EP 0139421;



(Id)


- 5 wherein R^1 , R^2 and R^3 are as defined in relation to formula (I) of EP 0032128;



(Ie)

10


- wherein A, R, R^1 and X are as defined in relation to formula (I) of EP 0428312;



(If)

15

- when A, B, R and R^1 are as defined in relation to formula (II) of EP 0428312;




(Ig)

20


- wherein R^1 is as defined in relation to formula (I) of EP 0489663;

- 5 -




(Ih)

wherein R^1 , R^2 , R^3 and n are as defined in relation to formula (I) of
 5 EP 0155845;




(Ii)

10 when R^1 is as defined in relation to formula (I) of EP 0257781;



(Ij)

15 wherein Ar, R^1 , R^2 , R^3 , R^4 , R^5 , n , U and W are as defined in relation to
 formula (I) of United States Patent No. 5104888;




(Ik)

20


when A, R^1 , R^2 and X are as defined in relation to formula (I) of
 EP 0208420;

- 6 -



(II)


- 5 when R¹, R², X, Z, m and n are as defined in relation to formula (I) of EP 0177353;



(Im)

10


according to formula (I) of EP 0319189;



(In)

15


wherein A, B, X, X¹, X², n and Z are as defined in relation to formula (I) of EP 0332331;



(Io)

20

wherein V, W, X, Y, Z, Z¹ and n are as defined in EP 0332332; and



(Ip)


25

wherein Q and X are as defined in relation to formula (I) of International Application No. WO 92/18501.


- 5 Favourably, T represents a moiety of the above defined formula (Ia), (Ic), (Ie), (If), (Ii), (Ik) or (Io).

In particular T represents a moiety selected from the list consisting of (a), (b), (c), (d), (e), (f), (g), (h) and (i):


10




(a)




(b)




(c)



(d)




(e)




(f)


15



(g)



(h), and




Suitably, T represents a moiety of formula (Ia) as defined above.

5 Preferably, T¹ represents S.

Thus, in a preferred aspect, the invention provides a process for preparing a compound of formula (I) as defined in EP 0306228: Accordingly, the invention provides a process for preparing a compound of formula (1):

10




15 or a tautomeric form thereof and/or a pharmaceutically acceptable salt thereof, and/or a pharmaceutically acceptable solvate thereof, wherein:
A¹ represents a substituted or unsubstituted aromatic heterocyclyl group;
R¹ represents a hydrogen atom, an alkyl group, an acyl group, an aralkyl group, wherein the aryl moiety may be substituted or unsubstituted, or a
20 substituted or unsubstituted aryl group;
A² represents a benzene ring having in total up to five substituents; and
n represents an integer in the range of from 2 to 6;

which process comprises, treating a compound of formula (2):

25

- 9 -



- or a tautomeric form thereof and/or a salt thereof, and/or a solvate thereof,
 5 wherein A^1 , A^2 , R^1 and n are as defined in relation to formula (1) with a microbial reductase obtained from an appropriate red yeast; and thereafter, as required, preparing a pharmaceutically acceptable salt, and/or a pharmaceutically acceptable solvate of the compound of formula (1) or a tautomeric form thereof.
- 10 Unless mentioned to the contrary herein, the suitable, apt, favoured and preferred values for each variable in the above mentioned moieties of formula (Ia), (Ib), (Ic), (Id), (Ie), (If), (Ig), (Ih), (Ii), (Ij), (Il), (Im), (In), (Io) or (Ip) are as defined in the European and International patent
 15 applications or United States patents mentioned above in respect of each of the said formulae.

In particular, the suitable, apt, favoured and preferred values of the variables A^1 , A^2 , R^1 and n in formula (1) and formula (2) are as defined
 20 in relation to formula (I) of EP 0306228.

A most preferred value of A^1 in formula (1) and in formula (2) is a 2-pyridyl group.

- 25 A most preferred value of A^2 in formula (1) and in formula (2) is a moiety of formula:



- 30 A most preferred value of R^1 in formula (1) and in formula (2) is a methyl group.

A most preferred value of n in formula (1) and in formula (2) is 2.

- 10 -

A most preferred value of T is a moiety of formula (a) as defined above.

A most preferred value of formula (1) is 5-(4-[2-(N-methyl-N-(2-pyridyl)amino)ethoxy]benzyl)-2,4-thiazolidinedione, or a tautomeric form thereof and/or a pharmaceutically acceptable salt thereof, and/or a pharmaceutically acceptable solvate thereof.

A most preferred value of formula (2) is 5-(4-[2-(N-methyl-N-(2-pyridyl)amino)ethoxy]benzylidene)-2,4-thiazolidinedione or a tautomeric form thereof and/or a salt thereof, and/or a solvate thereof.

Unless mentioned to the contrary herein, the suitable, apt, favoured and preferred pharmaceutically acceptable salts, pharmaceutically acceptable solvates and tautomeric forms of each of the compounds in the above mentioned moieties of formula (Ia), (Ib), (Ic), (Id), (Ie), (If), (Ig), (Ih), (Ii), (Ij), (Il), (Im), (In), (Io) or (Ip) are as defined in the European or International patent applications or United States patents mentioned above in respect of each of the said formulae.

In particular, suitable pharmaceutically acceptable salts include metal salts, such as for example aluminium, alkali metal salts such as sodium or potassium, alkaline earth metal salts such as calcium or magnesium and ammonium or substituted ammonium salts, for example those with lower alkylamines such as triethylamine, hydroxy alkylamines such as 2-hydroxyethylamine, bis-(2-hydroxyethyl)-amine or tri-(2-hydroxyethyl)-amine, cycloalkylamines such as bicyclohexylamine, or with procaine, dibenzylpiperidine, N-benzyl- β -phenethylamine, dehydroabietylamine, N,N'-bisdehydroabietylamine, glucamine, N-methylglucamine or bases of the pyridine type such as pyridine, collidine or quinoline.

In particular, suitable pharmaceutically acceptable solvates include hydrates.

An appropriate red yeast is a red yeast which provides the above mentioned reduction, including known red yeasts and those red yeasts which may be produced from known red yeasts by conventional methods, such as conventional mutation methods, including the use of ultra-violet light, chemical mutagenic agents and genetic manipulation.

Particular red yeasts include those species of the genera *Rhodotorula*, *Rhodospiridium* or synonyms thereof.

- 5 Particular examples of the genera *Rhodotorula* include *Rhodotorula glutinis* CBS 4406, *Rhodotorula rubra* CBS 6469, *Rhodotorula rubra* CBS 17 and *Rhodotorula glutinis* IFO 0869.

- 10 Particular examples of the genera *Rhodospiridium* include *Rhodospiridium toruloides* CBS 14.

Suitable species of the genera *Rhodotorula* and *Rhodospiridium* and synonyms thereof are as disclosed in 'The Yeasts', 2nd edition, 1970, J.Lodder (ed.), North-Holland Publishing Co.

- 15 Known red yeasts may be obtained from the appropriate collection such as those disclosed in the National Collection of Yeast Cultures Catalogue, 1990, AFRC, Institute of Food Research, Norwich, UK.

- 20 *Rhodotorula glutinis* CBS 4406, *Rhodotorula rubra* CBS 6469, *Rhodotorula rubra* CBS 17 and *Rhodospiridium toruloides* CBS 14 are known yeast microorganisms and may be obtained from Centraal bureau voor Schimmelcultures, Baarn, Delft, Netherlands. *Rhodotorula glutinis* IFO 0869 is a known yeast microorganism and may be obtained from the
25 Institute for Fermentation, Osaka, Japan.

- The compounds of formula (I) and formula (1) have at least one asymmetric carbon atom, indicated with an asterisk '*' in the formulae; they may therefore exist in at least two optically isomeric forms.

- 30 The compounds of formula (II) may be prepared according to known methods, for example by use of the appropriate method disclosed in the abovementioned European and International patent applications or U.S. patents. The contents of the abovementioned European and International
35 patent applications and U.S. patents are incorporated herein by reference.

In particular compounds of formula (2) may be prepared according to the methods disclosed in EP 0306228.

The microbial reductase enzymes may be obtained from the appropriate red yeast by conventional culturing techniques such as those disclosed in J. Bacteriol., 1982, Vol.150 498-505. H. Gilbert and M. Tully, European
5 Patent Application No. 0198440 and British Patent No. 1,474,519.

The reductase may be isolated as a pure enzyme or, in the alternative a suitable source of the reductase may be incorporated into the reaction.

10 Suitable sources of the reductase include whole yeast cell cultures or immobilised yeast cells from the abovementioned organisms, suitably wherein the yeast cells are *Rhodotorula glutinis* CBS 4406 or *Rhodotorula rubra* CBS 6469.

15 The organisms may be grown in any suitable growth medium, including yeast extract, synthetic medium, for example peptone, or mixtures thereof, for example a yeast extract / peptone broth, at any temperature commensurate with growth of the organism, being generally an ambient or slightly elevated temperature, such as a temperature in the range of
20 from 20°C to 50°C according to the nature of the organism employed, a suitable temperature being in the range of from 20°C to 40°C, favourably, suitably 20°C to 30°C, for example 28°C.

The reduction of the compound of formula (II) may be carried out in any
25 suitable solvent, including the above mentioned growth medium, or, after separation and transfer of the cells or immobilised cells, in a salt solution, such as a buffer, for example an aqueous citrate buffer pH3.75 containing 5 w/v sucrose.

30 Generally, the compound of formula (II) is introduced into the reaction system as a solution in an organic solvent which may be a water miscible solvent such as dioxan, a partially miscible solvent, such as methyl acetate, or a water immiscible solvent, such as ethyl acetoacetate. The reaction may then carried out in the resulting single or two phase system.


35 The reaction may be carried out at any pH which provides a suitable rate of formation of the required product which is generally a pH in the range of from 2 to 10, such as in the range of from 2 to 7, 4 to 8 or 7 to 10, or

- 13 -

more particularly in the range of from 2 to 4, 3 to 5, 4 to 6, 5 to 7, 6 to 8 or 8 to 10, for example at pH3 or pH8.

In one particular aspect of the invention, the reaction is carried out at an acidic pH, which is indicated to provide an stereoselective reduction of the substrate, to provide a product enriched in one enantiomer ('The selected enantiomer') of the asterisked carbon atom.

Accordingly, there is provided a process for the preparation of a compound of formula (I) (hereinafter referred to as the 'enantiomerically enriched compound (I)') wherein greater than 50% w/w of said compound is in the form of a compound of formula (IA) :



(IA)

or a tautomeric form thereof and/or a pharmaceutically acceptable salt thereof and/or a pharmaceutically acceptable solvate thereof, wherein T and T¹ are as defined in relation to formula (I) and the '*' carbon atom is an enantiomeric carbon atom, which process comprises reacting a compound of the above defined formula (II) with a microbial reductase obtained from an appropriate red yeast and wherein the reaction is carried out at an acidic pH; and thereafter, as required, preparing a pharmaceutically acceptable salt and/or a pharmaceutically acceptable solvate of the enantiomerically enriched compound (I) or a tautomeric form thereof.

The values of variable T and T¹ in the compound of formula (IA) are as discussed above in respect of the compound of formula (I).

A most preferred form of the compound of formula (IA) is (+) 5-(4-[2-(N-methyl-N-(2-pyridyl)amino)ethoxy]-benzyl)thiazolidine-2,4-dione, or a tautomeric form thereof and/or a pharmaceutically acceptable salt thereof and/or a pharmaceutically acceptable solvate thereof.

- 14 -

A suitable acidic pH is a pH in the range of from 2 to 6, especially 2 to 4, and preferably at pH3.

- The stereoselective process provides the enantiomerically enriched compound (I) enriched in the enantiomer having the same stereochemistry at the asterisked carbon atom as the equivalent carbon atom in (+) - 5-(4-[2-(N-methyl-N-(2-pyridyl)amino)ethoxy]-benzyl)thiazolidine-2,4-dione; X-ray analysis indicates this to be the (R)-enantiomer.
- 10 The reaction conditions, such as the particular acidic pH and the reaction temperature which provide optimum enrichment for any particular enantiomerically enriched compound (I) may be determined by routine experimentation.
- 15 Suitably, the stereoselective reaction provides enantiomerically enriched compound (I) wherein greater than 70% w/w is compound (IA); and favourably greater than 80% w/w. Most favourably, the product from the stereoselective process provides enantiomerically enriched compound (I) wherein 80-100% w/w is the compound of formula (IA), preferably 90-100%, such as 90-95%, and most preferably 95-100%, for example 95%, 96%, 97%, 98%, 99% or 100% w/w of compound of formula (IA).

- The above mentioned enantiomerically enriched compound (I) is considered to form a further aspect of the present invention. Accordingly the present invention provides enantiomerically enriched compound (I) or a tautomeric form thereof and/or a pharmaceutically acceptable salt thereof and/or a pharmaceutically acceptable solvate thereof.

- The present invention also provides enantiomerically enriched compound (I) or a tautomeric form thereof and/or a pharmaceutically acceptable salt thereof and/or a pharmaceutically acceptable solvate thereof, wherein greater than 50% w/w is in the form of compound (IA); suitably greater than 70% w/w and favourably greater than 80% w/w. Most favourably, the enantiomerically enriched compound (I) is in a form wherein 80-100% w/w is a compound of formula (IA), preferably 90-100%, such as 90-95%, and most preferably 95-100%, for example 95%, 96%, 97%, 98%, 99% or 100% w/w of a compound of formula (IA).

In one preferred aspect there is provided a compound of formula (IA) or a tautomeric form thereof and/or a pharmaceutically acceptable salt thereof and/or a pharmaceutically acceptable solvate thereof, preferably in optically pure form.

5

The absolute stereochemistry of compounds may be determined using conventional methods, such as X-ray crystallography.

10 The reductions may be carried out at any temperature which provides a suitable rate of formation of the required product, being generally an ambient or slightly elevated temperature, such as a temperature in the range of from 20°C to 50°C, such as in the range of from 20°C to 40°C and preferably in the range of from 20°C to 30°C, for example 28°C.

15 The product from any of the above reactions may be purified by phase separation and/or extraction into a suitable solvent, such as dichloromethane and thereafter, if required it may be chromatographed.

20 The above mentioned reactions may also be carried out using continuous production methods involving continuous recycling of the organic phase in the reaction or immobilisation of the cells and passing the substrate in organic solution over the immobilised biomass in a column, loop reactor or other similar reactor.

25 Immobilised cell preparations may be prepared according to conventional procedures, for example those disclosed in 'Alginate as immobilisation matrix for cells'; Smidsrod and Skjak-Braek, Tibtech 1990, 8, 71-78 or 'Immobilised enzymes and cells'; Rosevar, Kennedy and Cabral, IOP Publishing Ltd., 1987.

30

As mentioned above the enantiomerically enriched compound (I) is indicated as having useful therapeutic properties: The present invention accordingly provides enantiomerically enriched compound (I), or a tautomeric form thereof and/or a pharmaceutically acceptable salt thereof and/or a pharmaceutically acceptable solvate thereof, for use as an active
35 therapeutic substance.

Thus the present invention provides enantiomerically enriched compound (I), or a tautomeric form thereof and/or a pharmaceutically acceptable salt thereof and/or a pharmaceutically acceptable solvate thereof, for use in the treatment of and/or prophylaxis of hyperglycaemia.

5

In a further aspect the present invention also provides enantiomerically enriched compound (I), or a tautomeric form thereof and/or a pharmaceutically acceptable salt thereof and/or a pharmaceutically acceptable solvate thereof, for use in the treatment and/or prophylaxis of hyperlipidaemia, hypertension, cardiovascular disease and certain eating disorders.

10

Enantiomerically enriched compound (I), or a tautomeric form thereof and/or a pharmaceutically acceptable salt thereof and/or a pharmaceutically acceptable solvate thereof, may be administered per se or, preferably, as a pharmaceutical composition also comprising a pharmaceutically acceptable carrier.

15

Accordingly, the present invention also provides a pharmaceutical composition comprising enantiomerically enriched compound (I), or a tautomeric form thereof, or a pharmaceutically acceptable salt thereof, or a pharmaceutically acceptable solvate thereof, and a pharmaceutically acceptable carrier therefor.

20

As used herein the term 'pharmaceutically acceptable' embraces compounds, compositions and ingredients for both human and veterinary use: for example the term 'pharmaceutically acceptable salt' embraces a veterinarily acceptable salt.

25

The composition may, if desired, be in the form of a pack accompanied by written or printed instructions for use.

30

Usually the pharmaceutical compositions of the present invention will be adapted for oral administration, although compositions for administration by other routes, such as by injection and percutaneous absorption are also envisaged.

35

Particularly suitable compositions for oral administration are unit dosage forms such as tablets and capsules. Other fixed unit dosage forms, such as powders presented in sachets, may also be used.

- 5 In accordance with conventional pharmaceutical practice the carrier may comprise a diluent, filler, disintegrant, wetting agent, lubricant, colourant, flavourant or other conventional adjuvant.

- 10 Typical carriers include, for example, microcrystalline cellulose, starch, sodium starch glycollate, polyvinylpyrrolidone, polyvinylpolypyrrolidone, magnesium stearate or sodium lauryl sulphate.

- 15 Most suitably the composition will be formulated in unit dose form. Such unit dose will normally contain an amount of the active ingredient in the range of from 0.1 to 1000 mg, more usually 0.1 to 500 mg, and more especially 0.1 to 250 mg.

- 20 The present invention further provides a method for the treatment and/or prophylaxis of hyperglycaemia in a human or non-human mammal which comprises administering an effective, non-toxic, amount of enantiomerically enriched compound (I), or a tautomeric form thereof and/or a pharmaceutically acceptable salt thereof and/or a pharmaceutically acceptable solvate thereof to a hyperglycaemic human or non-human mammal in need thereof.

- 25 The present invention further provides a method for the treatment of hyperlipidaemia in a human or non-human mammal, which comprises administering an effective, non-toxic, amount of enantiomerically enriched compound (I), or a tautomeric form thereof and/or a pharmaceutically acceptable salt thereof and/or a pharmaceutically acceptable solvate thereof, to a hyperlipidaemic human or non-human mammal in need thereof.
- 30

- 35 Conveniently, the active ingredient may be administered as a pharmaceutical composition hereinbefore defined, and this forms a particular aspect of the present invention.

- 18 -

- In the treatment and/or prophylaxis of hyperglycaemic humans, and/or the treatment and/or prophylaxis of hyperlipidaemic human, the enantiomerically enriched compound (I), or a tautomeric form thereof and/or a pharmaceutically acceptable salt thereof and/or a pharmaceutically acceptable solvate thereof, may be taken in doses, such as those described above, one to six times a day in a manner such that the total daily dose for a 70 kg adult will generally be in the range of from 0.1 to 6000 mg, and more usually about 1 to 1500 mg.
- 10 In the treatment and/or prophylaxis of hyperglycaemic non-human mammals, especially dogs, the active ingredient may be administered by mouth, usually once or twice a day and in an amount in the range of from about 0.025 mg/kg to 25 mg/kg, for example 0.1 mg/kg to 20 mg/kg. Similar dosage regimens are suitable for the treatment and/or prophylaxis of hyperlipidaemia in non-human mammals.

The dosages regimens for the treatment of hypertension, cardiovascular disease and eating disorders will generally be those mentioned above in relation to hyperglycaemia.

- 20 In a further aspect the present invention provides the use of enantiomerically enriched compound (I), or a tautomeric form thereof and/or a pharmaceutically acceptable salt thereof and/or a pharmaceutically acceptable solvate thereof, for the manufacture of a medicament for the treatment and/or prophylaxis of hyperglycaemia.

- 30 The present invention also provides the use of enantiomerically enriched compound (I), or a tautomeric form thereof and/or a pharmaceutically acceptable salt thereof, and/or a pharmaceutically acceptable solvate thereof, for the manufacture of a medicament for the treatment and/or prophylaxis of hyperlipidaemia, hypertension, cardiovascular disease or certain eating disorders.

- 35 The following Examples illustrate the invention. It will be appreciated that catalytic activity, and hence yields may be improved by carrying out strain improvement on the organism, by known techniques.

Examples: Preparation of Microbiological Culture Medium

The following microbiological culture media were used in the examples given:

5

Medium A; yeast extract (10g), and mycological peptone (20g) were dissolved in de-ionised water to make 1 litre and the pH was adjusted to 7.0-7.2 by the addition of 2M sodium hydroxide solution. This was dispensed in 90ml volumes into 500ml Erlenmeyer flasks prior to sterilisation, after which 10 ml of 30% w/v D-glucose solution was added to each flask with filter sterilisation.

Medium B; $(\text{NH}_4)_2\text{HPO}_4$ (13g), KH_2PO_4 (7g), yeast extract (3g), $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$ (0.8g), NaCl (0.1g), $\text{ZnSO}_4 \cdot 7\text{H}_2\text{O}$ (60mg), $\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$ (90mg), $\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$ (5mg) and $\text{MnSO}_4 \cdot 4\text{H}_2\text{O}$ (10mg) were dissolved in de-ionised water to make 900ml and the pH adjusted to 7.0-7.2 by the addition of 2M sodium hydroxide solution. This was dispensed in 90ml aliquots into 500 ml Erlenmeyer flasks for pressure sterilisation after which 10ml of 40% (w/v) D-glucose solution was filter sterilised into each flask.

Example 1: Reduction Using Free Yeast Cells.

25 A loopful of *Rhodotorula rubra* CBS 6469 was used to inoculate a shake flask of medium A and this was incubated with shaking at 28°C for 72h. This broth (1ml) was used to inoculate a similar flask of medium A which was shaken as before for 48h prior to centrifugation. The pellet was resuspended in 0.1M Tris/HCl buffer pH8.0 containing 5% (w/v) sucrose giving a 1.3 relative broth cell density. To a portion of this suspension (40ml) in a 250ml Erlenmeyer flask was added 5-(4-[2-(N-methyl-N-(2-pyridyl)amino)ethoxy]benzylidene)thiazolidine-2,4-dione (7.5ml of a 5mg/ml solution in 1,4-dioxan) and the mixture shaken at 28°C for 22h. Centrifugation was followed by removal of 41.5ml of supernatant which by hplc indicated an 80% production of 5-(4-[2-(N-methyl-N-(2-pyridyl)amino)ethoxy]-benzyl)thiazolidine-2,4-dione.

Example 2: Reduction Using Immobilised Yeast Cells

A loopful of *Rhodotorula rubra* CBS 6469 was used to inoculate a shake flask of medium B and this was incubated with shaking at 28°C for 72h. This broth (1ml) was used to inoculate similar flasks of medium B which were shaken as before for 48h prior to centrifugation of 120ml of the broth. The resulting pellet was washed once in 0.1M Tris/HCl buffer pH8.0 and resuspended to 12.5ml in this buffer. An equal volume of 2% (w/v) sodium alginate solution in the same buffer was added and the cells immobilised using standard methodology (Alginate as immobilisation matrix for cells. Smidsrod and Skjak-Braek, Tibtech 1990, 8, 71-78). The resulting beads were washed in the Tris buffer previously described and the volume made up to 40ml with this buffer. To this suspension in a 250 ml Erlenmeyer flask was added 5-(4-[2-(N-methyl-N-(2-pyridyl)amino)ethoxy]benzylidene)thiazolidine-2,4-dione (7.5ml of a 5mg/ml solution in 1,4-dioxan) and the mixture shaken at 28°C for 22h. The supernatant was decanted and the beads washed with 50 ml of 20% (v/v) 1,4-dioxan in the Tris buffer. Hplc of the combined solutions indicated a conversion to 87% of 5-(4-[2-(N-methyl-N-(2-pyridyl)amino)ethoxy]benzyl)thiazolidine-2,4-dione. The product was extracted into dichloromethane and the organic phase dried over magnesium sulphate and evaporated to yield the product as confirmed by spectroscopy.

Example 3: Reduction Using Free Yeast Cells

A loopful of *Rhodotorula rubra* CBS 6469 was used to inoculate a shake flask of medium A and this was incubated with shaking at 28°C for 72h. This broth (1ml) was used to inoculate a similar flask of medium A which was shaken as before for 48h prior to centrifugation of 10ml of the broth. The pellet was resuspended in 0.3M Tris/HCl buffer pH8.0 containing 5% (w/v) sucrose to a volume of 4.0ml. To this suspension in a 25ml Erlenmeyer flask was added 5-(4-[2-(N-methyl-N-(2-pyridyl)amino)ethoxy]benzylidene)thiazolidine-2,4-dione (0.55ml of a 8.33mg/ml solution in 1,4-dioxan) and the mixture shaken at 28°C for 4h. Assay of the reaction mixture by hplc indicated a conversion to 59% to product (5-[4-(2-(N-methyl-N-(2-pyridyl)amino)ethoxy]-

- 21 -

benzyl)thiazolidine-2,4-dione) which had an enantiomer ratio identical to a racemic standard.

Example 4: Production of Enantiomerically Enhanced Product.

5

Rhodotorula rubra CBS 6469 was grown as described in Example 1, and the cell pellet from 66ml centrifuged medium was resuspended to 25ml in 0.1M citrate buffer containing 5% w/v sucrose at either pH 3.0, 3.5 or 4.0. To each was added 3.44ml of a 8.33mg/ml solution of 5-(4-[2-(N-methyl-N-(2-pyridyl)amino)ethoxy]benzylidene)thiazolidine-2,4-dione in 1,4-dioxan. After shaking in flasks for 4h the reaction mixtures were assayed by hplc for product.

10

pH	% conversion to product	enantiomer ratio of product
3.0	71	>98:<2
3.5	77	95:5
4.0	81	91:9

15 Using these results the pellet from 220ml broth was resuspended in 88ml of citrate buffer/sucrose pH 3.75 and to each of two 44ml portions in 500ml flasks was added 6ml of the above substrate solution. After 3h. 20min shaking at 28°C the reaction mixtures were centrifuged and the cells washed with 25ml of 12%v/v 1,4-dioxan in the above buffer system. Hplc indicated a 94:6 enantiomer ratio of product in the combined supernatant.

20

The solution was reduced to 2/3rds original volume under vacuum at room temperature to remove dioxan. The solution was basified to pH8 using 10% aqueous ammonia and extracted with dichloromethane (3 x 50ml).

25 The combined organic extracts were dried (MgSO₄), filtered, evaporated to dryness under vacuum at <25°C and the gummy product was dissolved in water (10ml) containing concentrated hydrochloric acid (0.2ml). After cooling to 2°C for 24 hours this solid was filtered and dried under vacuum (20°C) to give (+)-5-(4-[2-(N-methyl-N-(2-pyridyl)amino)ethoxy]-benzyl)thiazolidine-2,4-dione hydrochloride. (mp 123.5- 124°C; water)

30

Example 5: Reduction using *Rhodotorula glutinis* CBS 4406

Rhodotorula glutinis CBS 4406 was cultured as described in Example 2, to provide 120ml broth. The broth was centrifuged and resuspended to 12.5ml in pH 8.0, 0.1M tris/HCl buffer containing 5% (w/v) sucrose. An equal volume of 2% (w/v) sodium alginate was added and the cells were immobilised as in Example 2, then resuspended to 40ml in the above buffer. The suspension was dispensed into a 250ml Erlenmeyer flask, to which was added 7.5mls of a 5mg/ml solution of 5-(4-[2-(N-methyl-N-(2-pyridyl)amino)ethoxy]benzylidene)thiazolidine-2,4-dione in 1,4-dioxan. The flask was shaken for 22h at 28°C, then the supernatant was analysed using hplc which indicated a 51% production of 5-(4-[2-(N-methyl-N-(2-pyridyl)amino)ethoxy]benzyl)thiazolidine-2,4-dione.

Example 6: Reduction using *Rhodotorula rubra* CBS 6469


Rhodotorula rubra CBS 6469 was cultured as described in Example 1. Broth (12mls) was centrifuged and resuspended to 4.0ml in pH 9.0, 0.3M Tris/HCl buffer containing 5% (w/v) sucrose. The suspension was dispersed into a 25ml Erlenmeyer flask, to which was added 0.55ml of a 8.3mg/ml solution of 5-(4-[2-N-(2-benzoxazolyl)-(N-methyl)amino)ethoxy]benzylidene)thiazolidine-2,4-dione in 1,4-dioxan. The flask was shaken for 24h. at 28°C, then the broth was analysed using hplc which indicated a 49% production of 5-(4-[2-N-(2-benzoxazolyl)-(N-methyl)amino)ethoxy]benzyl)thiazolidine-2,4-dione.

30

35


Claims

1. A process for preparing a compound of formula (I):




(I)

- or a tautomeric form thereof and/or a pharmaceutically acceptable salt thereof and/or a pharmaceutically acceptable solvate thereof, wherein:
 10 T represents a substituted or unsubstituted aryl group and T¹ is O or S;
 which process comprises, treating a compound of formula (II):




(II)

- 15 or a tautomeric form thereof and/or a salt thereof and/or a solvate thereof,
 wherein T and T¹ are as defined in relation to formula (I), with a
 microbial reductase obtained from an appropriate red yeast; and
 thereafter, as required, preparing a pharmaceutically acceptable salt
 20 and/or a pharmaceutically acceptable solvate of the compound of formula
 (I) or a tautomeric form thereof.
2. A process according to claim 1, wherein T represents a moiety
 selected from the list consisting of (Ia), (Ib), (Ic), (Id), (Ie), (If), (Ig), (Ih),
 25 (Ii), (Ij), (Il), (Im), (In), (Io) and (Ip):




(Ia)

wherein A^1 , A^2 , R^1 and n are as defined in relation to formula (I) of EP
 5 0306228;




(Ib)

wherein R^2 , L^1 , L^2 and L^3 are as defined in relation to formula (I) of
 10 EP 0008203;



(Ic)


15 wherein R^1 , R^2 , R^3 , R^4 , R^5 , W and n are as defined in relation to formula
 (I) of EP 0139421;



(Id)


20 wherein R^1 , R^2 and R^3 are as defined in relation to formula (I) of
 EP 0032128;

-25-




(Ie)

wherein A, R, R¹ and X are as defined in relation to formula (I) of
 5 EP 0428312;



(If)


10 when A, B, R and R¹ are as defined in relation to formula (II) of
 EP 0428312;



(Ig)

15


wherein R¹ is as defined in relation to formula (I) of EP 0489663;



(Ih)

20


wherein R¹, R², R³ and n are as defined in relation to formula (I) of
 EP 0155845;



(Ii)

when R^1 is as defined in relation to formula (I) of EP 0257781;


5



(Ij)


wherein Ar, R^1 , R^2 , R^3 , R^4 , R^5 , n, U and W are as defined in relation to formula (I) of United States Patent No. 5104888;

10



(Ik)


15 when A, R^1 , R^2 and X are as defined in relation to formula (I) of EP 0208420;



(II)

20


when R^1 , R^2 , X, Z m and n are as defined in relation to formula (I) of EP 0177353;



(Im)


according to formula (I) of EP 0319189;

5




(In)

10 wherein A, B, X, X¹, X², n and Z are as defined in relation to formula (I) of EP 0332331;



(Io)

15 wherein V, W, X, Y, Z, Z¹ and n are as defined in EP 0332332; and




(Ip)


20 wherein Q and X are as defined in relation to formula (I) of International Application No. WO 92/18501.

3. A process according to claim 1 or claim 2, wherein T represents a moiety selected from the list consisting of (a), (b), (c), (d), (e), (f), (g), (h) and (i):


25




(a)




(b)




(c)



(d)




(e)




(f)


5



(g)



(h), and




(i)

4. A process according to any one of claims 1 to 3, wherein T represents a moiety of formula (Ia).

5. A process according to any one of claims 1 to 4, wherein T¹ represents S.

6. A process according to claim 1 for the preparation of a compound of formula (1):




(1)

- 10 or a tautomeric form thereof and/or a pharmaceutically acceptable salt thereof, and/or a pharmaceutically acceptable solvate thereof, wherein:
 A¹ represents a substituted or unsubstituted aromatic heterocyclyl group;
 R¹ represents a hydrogen atom, an alkyl group, an acyl group, an aralkyl group, wherein the aryl moiety may be substituted or unsubstituted, or a
 15 substituted or unsubstituted aryl group;
 A² represents a benzene ring having in total up to five substituents; and
 n represents an integer in the range of from 2 to 6;

which process comprises, treating a compound of formula (2):


20



(2)

- 25 or a tautomeric form thereof and/or a salt thereof, and/or a solvate thereof, wherein A¹, A², R¹ and n are as defined in relation to formula (1) with a microbial reductase obtained from an appropriate red yeast; and thereafter, as required, preparing a pharmaceutically acceptable salt, and/or a pharmaceutically acceptable solvate of the compound of formula (1) or a tautomeric form thereof.

7. A process according to claim 6, wherein the compound of formula (1) is 5-(4-[2-N-methyl-N-(2-pyridyl)amino]ethoxy]benzyl)-2,4-thiazolidinedione, or a tautomeric form thereof and/or a pharmaceutically acceptable salt thereof, and/or a pharmaceutically acceptable solvate thereof.
8. A process according to any one of claims 1 to 7, wherein an appropriate red yeast is a red yeast which provides the above mentioned reduction, including known red yeasts and those red yeasts which may be produced from known red yeasts by conventional methods.
9. A process according to any one of claims 1 to 8, wherein an appropriate red yeast is a red yeast from the species of the genera *Rhodotorula*, *Rhodospiridium* or synonyms thereof.
10. A process according to any one of claims 1 to 9 wherein an appropriate red yeast is *Rhodotorula glutinis* CBS 4406, *Rhodotorula rubra* CBS 6469, *Rhodotorula rubra* CBS 17 and *Rhodotorula glutinis* IFO 0869.
11. A process according to any one of claims 1 to 9 wherein an appropriate red yeast is *Rhodospiridium toruloides* CBS 14
12. A process for the preparation of a compound of formula (I) (the 'enantiomerically enriched compound (I)') wherein greater than 50% w/w of said compound is in the form of a compound of formula (IA) :



(IA)

30

or a tautomeric form thereof and/or a pharmaceutically acceptable salt thereof and/or a pharmaceutically acceptable solvate thereof, wherein T and T¹ are as defined in relation to formula (I) and the '*' carbon atom is an enantiomeric carbon atom, which process comprises reacting a

- compound of the above defined formula (II) with a microbial reductase obtained from an appropriate red yeast and wherein the reaction is carried out at an acidic pH; and thereafter, as required, preparing a pharmaceutically acceptable salt and/or a pharmaceutically acceptable solvate of the enantiomerically enriched compound (I) or a tautomeric form thereof.
13. Enantiomerically enriched compound (I) or a tautomeric form thereof and/or a pharmaceutically acceptable salt thereof and/or a pharmaceutically acceptable solvate thereof, wherein greater than 50% w/w is in the form of compound (IA)
14. A compound of formula (IA) or a tautomeric form thereof and/or a pharmaceutically acceptable salt thereof and/or a pharmaceutically acceptable solvate thereof, in optically pure form.
15. Enantiomerically enriched compound (I), or a tautomeric form thereof and/or a pharmaceutically acceptable salt thereof and/or a pharmaceutically acceptable solvate thereof, for use as an active therapeutic substance.
16. Enantiomerically enriched compound (I), or a tautomeric form thereof and/or a pharmaceutically acceptable salt thereof and/or a pharmaceutically acceptable solvate thereof, for use in the treatment of and/or prophylaxis of hyperglycaemia, hyperlipidaemia, hypertension, cardiovascular disease and certain eating disorders.
17. A pharmaceutical composition comprising enantiomerically enriched compound (I), or a tautomeric form thereof, or a pharmaceutically acceptable salt thereof, or a pharmaceutically acceptable solvate thereof, and a pharmaceutically acceptable carrier therefor.

INTERNATIONAL SEARCH REPORT

PCT/GB 92/02140

International Application No

I. CLASSIFICATION OF SUBJECT MATTER (If several classification symbols apply, indicate all) ⁶		
According to International Patent Classification (IPC) or to both National Classification and IPC		
Int.Cl. 5 C12P17/14; C07D413/12;	C07D277/34; A61K31/425;	C07D263/44; A61K31/42 C07D417/12
II. FIELDS SEARCHED		
Minimum Documentation Searched ⁷		
Classification System	Classification Symbols	
Int.Cl. 5	C12P ; C07D ; A61K	
Documentation Searched other than Minimum Documentation to the extent that such Documents are included in the Fields Searched ⁸		
III. DOCUMENTS CONSIDERED TO BE RELEVANT⁹		
Category ¹⁰	Citation of Document, ¹¹ with indication, where appropriate, of the relevant passages ¹²	Relevant to Claims No. ¹³
X	CHEMICAL AND PHARMACEUTICAL BULLETIN. vol. 32, no. 9, 1984, TOKYO JP pages 4460 - 4465 TAKASHI SOHDA ET AL. 'Studies on antidiabetic agents.VI. Asymmetric transformation of (+/-)-5-[4-(1-Methylcyclohexylmethoxy)benzyl]-2,4-thiazolidinedione (Ciglitazone) with optically active 1-phenylethylamines.' see the whole document	13-17
A	EP,A,0 306 228 (BEECHAM) 8 March 1989 cited in the application see examples 3-5 see examples 28-30 see claims	1
<p>¹⁰ Special categories of cited documents : ¹⁰</p> <p>"A" document defining the general state of the art which is not considered to be of particular relevance</p> <p>"E" earlier document but published on or after the international filing date</p> <p>"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>"O" document referring to an oral disclosure, use, exhibition or other means</p> <p>"P" document published prior to the international filing date but later than the priority date claimed</p> <p>"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step</p> <p>"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.</p> <p>"A" document member of the same patent family</p>		
IV. CERTIFICATION		
Date of the Actual Completion of the International Search	Date of Mailing of this International Search Report	
03 MARCH 1993	18. 03. 93	
International Searching Authority	Signature of Authorized Officer	
EUR PEAN PATENT OFFICE	DELANGHE L.L.M.	

III. DOCUMENTS CONSIDERED TO BE RELEVANT (CONTINUED FROM THE SECOND SHEET)		
Category *	Citation of Document, with indication, where appropriate, of the relevant passages	Relevant to Claim No.
A	EP,A,0 008 203 (TAKEDA) 20 February 1980 cited in the application see claims	1
A	EP,A,0 198 440 (KANEKAFUCHI) 22 October 1986 cited in the application see claims	1

**ANNEX TO THE INTERNATIONAL SEARCH REPORT
ON INTERNATIONAL PATENT APPLICATION NO.**

GB 9202140
SA 67612

This annex lists the patent family members relating to the patent documents cited in the above-mentioned international search report. The members are as contained in the European Patent Office EDP file on
The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

03/03/93

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP-A-0306228	08-03-89	AU-A- 2173888	09-03-89
		JP-A- 1131169	24-05-89
		US-A- 5002953	26-03-91
EP-A-0008203	20-02-80	JP-C- 1433701	07-04-88
		JP-A- 55022636	18-02-80
		JP-B- 62042903	10-09-87
		CA-A- 1131644	14-09-82
		US-A- 4287200	01-09-81
		US-A- 4340605	20-07-82
		US-A- 4438141	20-03-84
		US-A- 4444779	24-04-84
EP-A-0198440	22-10-86	JP-A- 61239893	25-10-86
		JP-A- 62065686	24-03-87
		JP-A- 62065687	24-03-87
		DE-A- 3686502	01-10-92
		JP-B- 4068915	04-11-92
		JP-A- 62029998	07-02-87
		US-A- 4857468	15-08-89